







Set up and implementation of a platform for the production and quality control of biological drugs













WP2 Tasks

Task 2.1: Facility implementation for process development and the production of preclinical lots of biologics


Task 2.2: Quality control laboratory for biologics and *in vitro* testing

Task 2.3: Animal facility of selected mouse models for *in vivo* efficacy studies









Guanidinoacetate methyltransferase (**GAMT**)


Methionine adenosyltransferase (**MAT2A**)

Phenylalanine ammonia-lyase (PAL)

Human α-mannosidase

Ubiquitin-specific protease 2 catalytic domain (Usp2)

mini-Cas9 (sRGN)



Recombinant Antibodies (Abs)

PROTEINS

Recombinant

Enzymes

Full-length (Dia-T51, anti-β-1,3 glucans)


scFv (single chain-fragment variable):

scFv-3T (anti-β-1,3 glucans); anti-ICAM-1; anti-IGF-1; anti-INSR

Conventional antibody

IVT modified mRNA

Guanidinoacetate methyltransferase (**GAMT**) mRNA
Ataxia Telangiectasia Mutated (**ATM**) mRNA









Recombinant protein expression workflow

Select/Design the end product (amino acid sequence)

Choose expression system



Design expression clone (DNA construct)




Express the protein



Purify the protein



Characterize the protein









Recombinant proteins generated


	Trecombinant proteins Senerated								
	Bacteria	Yeast	Plants						
Successfully produced *	 GAMT * MAT2A PAL* USP2* sRGN UBB* 	• MAT2A • PAL*	• α-Mannosidase*						
	 Ub-GFP* Dia-T51* scFv-3T* scFv anti-IGF1* scFv anti-INSR* scFv anti-ICAM-1* 								








Phenylalanine Ammonia Lyase (PAL) expression in Bacteria








3

Activity Assay by Mass Spectrometry










Anti-ICAM-1 scFv expression in Bacteria











PAL and MAT2A expression in Yeast











Ongoin:

Patent application Research article

Nico Betterle.

Biotechnology, University of Verona, Italy.

Plant-based biotechnology platform for a sustainable production of human recombinant α-mannosidase



A rare genetic disorder caused by a

deficiency of a human lysosomal enzyme.


It affects approximately

1 in 500,000 living births.



Plant as bioreactors (N.tabacum) for recombinant human enzyme production








Development of new innovative biosystems for a sustainable production

Vertical Farming – Hydroponic culture system



Kristina Ljumovic Deparment of

Enzymatic assay demonstrates promising results for future therapeutic applications

	Enzyme	
Leaves sample	Units/mg	
Wild-type	0.046	
Young transgenic leaves	0.47	10x 30x
Medium transgenic leaves	0.66	
Old transgenic leaves	1.327	







Ongoin:

Patent application Research article

Plant-based biotechnology platform for a sustainable production of human recombinant α-mannosidase













A rare genetic disorder caused by a deficiency of a human lysosomal enzyme. It affects approximately

1 in 500,000 living births.

Biotechnology, University of Verona, Italy.

Plant as bioreactors
(N.tabacum)
for recombinant human
enzyme production

Culture system optimization (Vertical Farming)

to improve scalability

innovative biosystems for a sustainable production

Development of new

Enzymatic assay demonstrates promising results for future therapeutic applications

Vertical Farming – Hydroponic culture system




	Enzyme	
Leaves sample	Units/mg	_
Wild-type	0.046	
Young transgenic leaves	0.47	10x 30x
Medium transgenic leaves	0.66	
Old transgenic leaves	1.327	









QC TESTS: BIOPHYSICAL CHARACTERIZATION OF RECOMBINANT PRODUCTS: the example of full-length antibody Dia-



QC Lab: some of these instruments were purchased by PNRR VITALITY projects




















CELL CYCLE

300 -

200 -

100 -

DMSO (ctrl)

%G1


44,6


%S


44,0

%G2

9,78









CELL VIABILITY ASSAY IN CELL LINES OF BOTH HEMATOPOIETIC AND SOLID

	TILLABA	
Cell line	IC50 (24h)	IC50 (48h)
U937	0.76	0.65
K562	1.25	1.18
Jurkat	0.73	0.28
NB4	0.50	0.37
U373	6.67	1.62
HeLa	1.67	0.94

SEARCHING FOR MOLECULAR MECHANISMS

WB ANALYSIS of Histopes PTMs and cMyc





	DMSO	Ak 12_0.5uM
G1 % (mean)	44,5 ± 1,7%	43,8 ± 1,3%
S % (mean)	43,8 ± 1,5%	40,6 ± 1,2%
G2 % (mean)	9,2 ± 1,4%	14,1 ± 1%







Monoclonal Antibody Conjugation with stable metal complex

GOALS?


Targeted Delivery:


Antibodies to be used as specific delivery vehicles to deliver metal complexes (e.g., radiometals) to target sites like tumors, increasing efficacy and reducing side effects.


Imaging:

Radiometals conjugated to antibodies are used in diagnostic imaging techniques to visualize specific tissues or cells.











By carrying a therapeutic radionuclide or cytotoxic drug, the antibody-metal complex can deliver radiation or a drug directly to diseased cells.











% Survival

100

100

100

100


0


57.1


42.9



In vivo studies of the antifungal humanized monoclonal antibody DIA-T51 ...







		Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
	C. auris	Lot of	5	737	NEW YEAR	RY.	13	
atment	Dia-T51 1 mg/kg + <i>C. auri</i> s	The	(3)	13			37	17
Pre-treatment	Dia-T51 1 mg/kg + <i>C. auris</i> + Dia-T51 1 mg/kg	3		6	52)	3 4	31	7





% Survival

100

100

100

100


0


57.1


42.9



In vivo studies of the antifungal humanized monoclonal antibody DIA-T51 ...








		Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
	C. auris	24.4	£43	13h	(C)	RY)	12	
Pre-treatment	Dia-T51 1 mg/kg + <i>C. auris</i>	The	(3)	13			(LE)	
	Dia-T51 1 mg/kg + <i>C. auris</i> + Dia-T51 1 mg/kg	0		8-	57	3 4	31	3










In vivo studies of the antifungal humanized monoclonal antibody DIA-T51 ...








... and its synergy with AMPHOTERICIN B






		Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
	C. auris	24.4	5	737	NEW YORK	R.	13	3
Pre-treatment	Dia-T51 1 mg/kg + <i>C. auri</i> s	The same	(3)				33	
	Dia-T51 1 mg/kg + <i>C. auris</i> + Dia-T51 1 mg/kg	0		6	47	3 4	51	2







Modified mRNA to treat monogenic diseases




GAMT mRNA

- IVT-modified mRNA
- Mouse model of GAMT deficiency for in vivo studies



ATM mRNA

- CDS for ATM variants cloned in pMRNAxp vector
- ATM-deficient cell lines for in vitro studies










RNA-based therapy for GAMT deficiency: in vitro studies



GAMT mRNA lipofection in HepG2

QPCR analysis

1,0

47

0,8

0,5

0,0

24h


48h

4

WB analysis

carrying IVT GAMT mRNA

- 1: Transfected HepG2 24h
- 2: Transfected HepG2 48h
- 3: Control HepG2 24h
- 4: Control HepG2 48h










RNA-based therapy for GAMT deficiency: in vivo studies











mRNA-based therapy for ATM deficiency: in vitro studies





































Thank you for your attention!